

A Note on Invariant Measure on the Local Gauge Group

Wei-Min Sun* Xiang-Song Chen and Fan Wang†
Department of Physics and Center for Theoretical Physics
Nanjing University, Nanjing 210093, China

Abstract

In this paper we investigate the problem of the existence of invariant measures on the local gauge group. We prove that it is impossible to define a *finite* translationally invariant measure on the local gauge group $C^\infty(\mathbf{R}^n, G)$ (where G is an arbitrary matrix Lie group).

Functional integral over the local gauge group is a commonly employed technique in the study of gauge field theories. Its earliest appearance in the literature is Faddeev and Popov's paper in 1967 [1], which investigated the quantization of non-Abelian gauge fields. In [2] this technique was used to construct a projection operator which implements the Gauss law, and in [3] it was used to construct a gauge invariant ground state wave-functional. In all these applications a key point is the assumption that we can define a translationally invariant measure on the local gauge group. It is known that on locally compact groups this is possible [4]. On non-locally-compact groups we cannot guarantee such a measure exists. In the case of vector spaces a standard result [5] says that an invariant measure exists only when the vector space is finite dimensional, in which case it is locally compact. The local gauge group is infinite dimensional and not locally compact. Since functional integral over the local gauge group is an important technique the existence of an invariant measure deserves a careful study. In [6] it is proved that on the local gauge group $C^\infty(\mathbf{R}^n, G)$ there exists no finite translationally invariant Borel measure (Haar measure). In this paper we study the case of the local gauge group $C^\infty(\mathbf{R}^n, G)$ (where G is an arbitrary matrix Lie group) and prove that it is impossible to define a finite translationally invariant Borel measure on it (provided that it is endowed with a suitable topology).

The local gauge group $G_0 = C^\infty(\mathbf{R}^n, G)$ is the group of all smooth mappings from $\mathbf{R}^n$ to G . For a generic element $\omega(x^1, \dots, x^n)$ of G_0 , the quantity $\omega^{-1}(0)\partial_1\omega(0)$ (here by $\partial_1\omega(0)$ we certainly mean $\frac{\partial}{\partial x^1}\omega(x)|_{x=0}$) belongs to the Lie algebra g of G . For $\xi = \sum_{i=1}^m u_i \xi_i \in g$ (where $\{\xi_1 \dots \xi_m\}$ is a basis of g) we define $f(\xi) = e^{-\sum_{i=1}^m u_i^2}$ and consider the following bounded continuous function of ω (in some suitable topology of the group G_0):

$$F[\omega] = f(\omega^{-1}(0)\partial_1\omega(0)) \quad (1)$$

*e-mail address: sunwm@chenwang.nju.edu.cn

†e-mail address: fgwang@chenwang.nju.edu.cn

In the following we shall prove that it is impossible to define a *finite* (left or right) translationally invariant measure on G_0 by using the method of *reduction to absurdity*. Suppose we have defined a measure $D\mu(\omega)$ which is right translationally invariant. Then we have the following equality:

$$\int_{G_0} D\mu(\omega) F[\omega] = \int_{G_0} D\mu(\omega) F[\omega\omega_0], \forall \omega_0 \in G_0 \quad (2)$$

From Eq(1) we have

$$\begin{aligned} & \int_{G_0} D\mu(\omega) f(\omega^{-1}(0)\partial_1\omega(0)) \\ &= \int_{G_0} f(\omega_0^{-1}(0)\omega^{-1}(0)\partial_1\omega(0)\omega_0(0) + \omega_0^{-1}(0)\partial_1\omega_0(0)), \forall \omega_0 \in G_0 \end{aligned} \quad (3)$$

In Eq(3) we set $\omega_0(x^1, \dots, x^n) = e^{\xi x^1}$, where ξ is an arbitrary element of the Lie algebra g and get

$$\int_{G_0} D\mu(\omega) f(\omega^{-1}(0)\partial_1\omega(0)) = \int_{G_0} D\mu(\omega) f(\omega^{-1}(0)\partial_1\omega(0) + \xi), \forall \xi \in g \quad (4)$$

Let us consider the continuous function $f(\omega^{-1}(0)\partial_1\omega(0) + \xi)$ defined on $G_0 \times g$ and observe that $(dm(\xi) = du^1 \dots du^m)$ is the Lebesgue measure on g

$$\begin{aligned} \int_{G_0} D\mu(\omega) \int_g dm(\xi) |f(\omega^{-1}(0)\partial_1\omega(0) + \xi)| &= \int_{G_0} D\mu(\omega) \int_g dm(\xi) f(\omega^{-1}(0)\partial_1\omega(0) + \xi) \\ &= \int_{G_0} D\mu(\omega) \int_g dm(\xi) f(\xi) \\ &= \pi^{n/2} \mu(G_0) \end{aligned} \quad (5)$$

where we have used the translational invariance of the Lebesgue measure $dm(\xi)$. According to Fubini theorem another iterated integral also exists and is equal to the integral in Eq(5):

$$\int_g dm(\xi) \int_{G_0} D\mu(\omega) f(\omega^{-1}(0)\partial_1\omega(0) + \xi) = \pi^{n/2} \mu(G_0) \quad (6)$$

However, according to Eq(4), which expresses the right translational invariance of $D\mu(\omega)$, we ought to have

$$\int_g dm(\xi) \int_{G_0} D\mu(\omega) f(\omega^{-1}(0)\partial_1\omega(0) + \xi) = +\infty \quad (7)$$

(we have $\int_{G_0} D\mu(\omega) f(\omega^{-1}(0)\partial_1\omega(0)) > 0$). Thus we get a contradiction, which shows that a *finite* right translationally invariant measure cannot be defined on G_0 . By considering the function $f(\omega(0)\partial_1\omega^{-1}(0))$ we can similarly show that a *finite* left translationally invariant measure cannot be defined on G_0 either.

The result proved above does not put essential constraints to the Faddeev-Popov formalism. It only requires that the volume of the local gauge group not be finite, which is a commonly-taken understanding. It also does not affect the construction of the gauge invariant ground state wave-functional described in [3]. However the construction of the

projection operator for the Gauss law described in [2] does have difficulties. In this reference the invariant measure is formally taken to be $D\mu(\omega) = \prod_x d\mu(\omega(x))$ where $d\mu(\omega(x))$ is the usual invariant measure of group G attached to the point x normalized to be $\int d\mu(\omega(x)) = 1$. However, the formal product of the measure $d\mu(\omega(x))$ over the whole space $\mathbf{R}^n$ is not a measure in the strict mathematical sense, and a finite invariant measure does not exist on the local gauge group.

REFERENCES

- [1] L.Faddeev and V.N.Popov, Phys.Lett. B25(1967),29.
- [2] Kerson Huang, *Quarks, Leptons and Gauge Fields* 2nd ed. World Scientific, 1992, sect.8.8.
- [3] I.I.Kogan and A.Kovner, Phys.Rev.D51(1995),1948.
- [4] P.R.Halmos, *Measure Theory*, Graduate Texts in Mathematics 18, Springer Verlag, 1974.
- [5] Y.Yamasaki, *Measures on Infinite Dimensional Spaces*, World Scientific, 1985.
- [6] W.M.Sun, Xiang-Song Chen and Fan Wang, hep-th/0105150